

*C2*

6. A process for the extraction of glycomacropeptide or caseinoglycomacropeptide ("GMP") from a lactic raw material comprising the steps of:

deionizing a lactic raw material for a time sufficient to obtain a substantially deionized lactic raw material having a pH of about 1 to 4.5 with the pH being adjusted, if necessary, to the recited range;

*start D3*

contacting the substantially deionized lactic raw material with an anionic resin having a hydrophobic matrix for a sufficient amount of time and at a sufficient temperature to remove GMP from the substantially deionized lactic raw material and to obtain a treated liquid material, wherein the substantially deionized lactic raw material contacts the resin in a gently stirred reactor at a temperature of less than 50°C for one to ten hours to adsorb the GMP onto the resin;

separating the resin from the treated liquid material; and  
rinsing the resin to obtain the GMP therefrom.

*C3*

9. A process for the extraction and removal of glycomacropeptide or caseinoglycomacropeptide ("GMP") from a lactic raw material comprising the steps of:

deionizing a lactic raw material for a time sufficient to obtain a substantially deionized lactic raw material having a pH of about 1 to 4.5 with the pH being adjusted, if necessary, to the recited range;

*start D4*

contacting the substantially deionized lactic raw material with an anionic resin having a hydrophobic matrix for a sufficient amount of time and at a sufficient temperature to remove GMP from the substantially deionized lactic raw material and to obtain a treated liquid material;

separating the resin from the treated liquid material; and  
concentrating the treated liquid material by evaporation and drying.

*C4 sub E1*

11. The process according to claim 1 wherein the anionic resin and the deionized lactic raw material are present in a ratio by volume of between 1:1 and 1:30.

*C5 sub E1*

14. The process of claim 1 wherein the treated liquid material has an amino acid profile is reduced in threonine and enriched in aromatic amino acids and tryptophan relative to the lactic raw material.